

INSTRUCTIONS FOLLOWING ORAL SURGERY

Brantford Central Dental www.brantfordcentraldental.com
Email: sheerdentalcare@gmail.com Phone: 519-759-7631

1. Bite on gauze for 1/2 hour.
2. Keep fingers and tongue away from the operated area.
3. Do not rinse your mouth until tomorrow, although it is permissible to drink cool or lukewarm liquids. Smoking should be avoided. Do not spit and use a straw if possible.
4. Starting tomorrow, rinse your mouth frequently with a solution of 1/2 teaspoon of salt in a glass of warm water. Continue the rinses for a few days.
5. If you have any discomfort, take whatever you normally take for a headache, every few hours. If necessary, the doctor will prescribe stronger medication for you.
6. Diet--Cold or lukewarm liquids may be taken for the first 4-6 hours. After this any soft food is permissible.
7. Bleeding--it is normal for the saliva to be streaked with blood for a day. If frank bleeding is present, fold sterile gauze into a firm wad and place it directly on the bleeding area. Maintain firm pressure by biting for 20 minutes. The gauze may be substituted by a tea bag. The tannic acid in tea has a clotting effect.
8. Swelling and Discoloration--is to be expected in certain areas, usually reaching its maximum two days after surgery. It will disappear gradually and is no cause for concern. If desired, an ice pad may be applied for the first 4-6 hours only, alternating 20 minutes on, 20 minutes off.
9. Sutures (Stitches)--if required, are removed without discomfort in about 5 days. An appointment will be made for you.
10. Do not hesitate to call the office if in doubt. If you feel the need to contact the dentist after hours, her emergency number is on our voice mail system.

INSTRUCTIONS FOLLOWING ORAL SURGERY

Brantford Central Dental www.brantfordcentraldental.com
Email: sheerdentalcare@gmail.com Phone: 519-759-7631

1. Bite on gauze for 1/2 hour.
2. Keep fingers and tongue away from the operated area.
3. Do not rinse your mouth until tomorrow, although it is permissible to drink cool or lukewarm liquids. Smoking should be avoided. Do not spit and use a straw if possible.
4. Starting tomorrow, rinse your mouth frequently with a solution of 1/2 teaspoon of salt in a glass of warm water. Continue the rinses for a few days.
5. If you have any discomfort, take whatever you normally take for a headache, every few hours. If necessary, the doctor will prescribe stronger medication for you.
6. Diet--Cold or lukewarm liquids may be taken for the first 4-6 hours. After this any soft food is permissible.
7. Bleeding--it is normal for the saliva to be streaked with blood for a day. If frank bleeding is present, fold sterile gauze into a firm wad and place it directly on the bleeding area. Maintain firm pressure by biting for 20 minutes. The gauze may be substituted by a tea bag. The tannic acid in tea has a clotting effect.
8. Swelling and Discoloration--is to be expected in certain areas, usually reaching its maximum two days after surgery. It will disappear gradually and is no cause for concern. If desired, an ice pad may be applied for the first 4-6 hours only, alternating 20 minutes on, 20 minutes off.
9. Sutures (Stitches)--if required, are removed without discomfort in about 5 days. An appointment will be made for you.
10. Do not hesitate to call the office if in doubt. If you feel the need to contact the dentist after hours, her emergency number is on our voice mail system.